

# Choice of Pore Size Can Introduce Artefacts when Filtering Picoeukaryotes for Molecular Biodiversity Studies

Nikolaj Sørensen · Niels Daugbjerg ·  
Katherine Richardson

Received: 26 October 2012 / Accepted: 26 December 2012 / Published online: 17 January 2013  
© Springer Science+Business Media New York 2013

**Abstract** Published results of studies based on samples size fractionated by sequential filtration (e.g. 0.2–3  $\mu\text{m}$ ) indicate that many ciliate, dinoflagellate and rhizarian phylotypes are found among marine picoeukaryotes. This is somewhat surprising as these protists are typically known as being large organisms (often  $>10 \mu\text{m}$ ) and no picoplanktonic species have so far been identified. Here, the abundances of ciliate and dinoflagellate phylotypes in published molecular studies of picoeukaryotes are shown to correlate negatively with the pore size chosen for the end filter in the sequential filtrations (i.e. the filter used to collect the microbial biomass). This suggests that extracellular DNA adhering to small particles may be the source of ciliate and dinoflagellate phylotypes in picoplanktonic size fractions. This hypothesis was confirmed using real-time qPCR, which revealed significantly less dinoflagellate 18S rDNA in a 0.8–3- $\mu\text{m}$  size fraction compared to 0.2–3  $\mu\text{m}$ . On average, the abundance of putative extracellular phylotypes decreased by 84–89 % when a 0.8- $\mu\text{m}$  end filter was used rather than a 0.2- $\mu\text{m}$  end filter. A 0.8- $\mu\text{m}$  filter is, however, not sufficient to retain all picoeukaryotic cells. Thus, selection of filter pore size involves a trade-off between avoiding artefacts generated by extracellular DNA and sampling the entire picoeukaryotic community. In contrast to ciliate and dinoflagellate phylotypes, rhizarian phylotypes in the picoplankton

size range do not display a pattern consistent with an extracellular origin. This is likely due to the documented existence of picoplanktonic swarmer cells within this group.

## Introduction

The first environmental molecular studies of marine picoeukaryotes (commonly defined as  $<3 \mu\text{m}$ ) were published in 2001 [9, 18, 29]. Since then, many more have followed and the increasing availability of high-throughput sequencing increases the likelihood that we will see many more such studies in the near future. Before embarking on such studies, it is important that all information relevant to their planning be extracted from the existing literature. One finding from the available literature that deserves some further consideration is the apparent abundance, diversity and ubiquity of ciliates, dinoflagellates and rhizarians found among the picoplankton [26, 39]. Unlike the diverse clades of marine stramenopiles (MAST) [22, 27] and the novel phylum of picobiliphytes [33], microscopic confirmation of these organisms within the picoplanktonic size fraction has not been obtained. The known examples of these groups are usually relatively large (commonly  $>10 \mu\text{m}$ ). It has earlier been suggested that extracellular (detrital) DNA from larger organisms may explain the apparent occurrence of these phylotypes among the picoplankton [23, 34]. If this is the case, we may be developing an incorrect understanding of biodiversity among the picoplankton.

Here, a systematic meta-analysis is made of picoeukaryotic environmental molecular studies in an attempt to identify potential relationships between the pore size of the end filter used when size fractioning picoeukaryotic samples by filtration and the abundance of ciliate, dinoflagellate and rhizarian DNA sequences. The meta-analysis indicated that less DNA is retained on filters with larger pore sizes compared to those having smaller ones. This suggests that much

**Electronic supplementary material** The online version of this article (doi:10.1007/s00248-012-0174-z) contains supplementary material, which is available to authorized users.

N. Sørensen · K. Richardson  
Center for Macroecology, Evolution and Climate,  
University of Copenhagen, Universitetsparken 15,  
2100 Copenhagen, Denmark

N. Sørensen (✉) · N. Daugbjerg  
Marine Biological Section, Department of Biology,  
University of Copenhagen, Universitetsparken 4,  
2100 Copenhagen, Denmark  
e-mail: nsorensen@bio.ku.dk

of the DNA recorded in such studies may be of extracellular origin. This hypothesis was subsequently tested experimentally using novel dinoflagellate-specific primers in a real-time qPCR set-up. The ability of different end filters to retain *Ostreococcus tauri*, the smallest known eukaryote (cell diameter of 0.8  $\mu\text{m}$  [6]), was also examined in this study.

## Methods

### Literature Analysis

To test for a correlation between the pore size of the end filter used and the abundance of ciliate, dinoflagellate and rhizarian DNA, relevant data were extracted from all published studies on the molecular diversity of picoeukaryotes found in the literature (Table S1). The abundances of dinoflagellates, ciliates and rhizarians in the collected data were calculated as the relative proportion of the total number of protist clones/tags (i.e. eukaryotes excluding fungi and metazoans). For comparative reasons, studies that presented diversity in a number of operational taxonomic units rather than clones were not included in the statistical analysis. Likewise, studies that used GF/F as the end filter were excluded, as the fibrous nature of glass fibre filters makes their pore size incomparable with those of other types of filters (e.g. Nuclepore).

### Real-Time qPCR Experiments

Samples for real-time qPCR were collected in the Fram Strait (78.782°N, 3.116°E) on the 9th of September 2011 at a 15-m depth using a rosette of 10-L Niskin bottles on a cruise with the R/V Lance. Under a gentle vacuum established using a hand pump, a volume of 1 L water was pre-filtered onto a 3- $\mu\text{m}$  filter and the microbial biomass was collected on either a 0.2- or 0.8- $\mu\text{m}$  filter (Nuclepore polycarbonate membranes, Whatman), for three and five samples, respectively. Filters were cut in half and stored at  $-20\text{ }^{\circ}\text{C}$  until DNA extraction. Only one-half of each filter was used for subsequent analysis.

Using 1.5-mL Eppendorf tubes, filters were incubated in 594  $\mu\text{L}$  CTAB and 6  $\mu\text{L}$   $\beta$ -mercaptoethanol at  $65\text{ }^{\circ}\text{C}$  and shaken at 1,000 rpm for 45 min on a PHMT Thermoshaker (Grant Instruments). Samples were then frozen for 30 min at  $-80\text{ }^{\circ}\text{C}$ , reheated at  $65\text{ }^{\circ}\text{C}$  for 30 min at 1,000 rpm and then heated for an additional 15 min without shaking. After adding 500  $\mu\text{L}$  of chloroform:isoamyl alcohol (volume 24:1), the samples were shaken continuously and vortexed twice for 10 min and then centrifuged for 5 min. The water phases were transferred to new Eppendorf tubes, and the procedure with chloroform:isoamyl alcohol was repeated. Two-thirds

of the sample volume of ice-cold isopropanol was then added, and the samples were kept at  $-20\text{ }^{\circ}\text{C}$  for 30 min. The samples were centrifuged for 10 min and washed in 500  $\mu\text{L}$  of 70 % ethanol and 50  $\mu\text{L}$  of 3 M Na-acetate buffer, centrifuged for 2 min and washed in 1,000  $\mu\text{L}$  96 % ethanol. Residual ethanol was evaporated by putting the opened sample tube on a heating block at  $65\text{ }^{\circ}\text{C}$  for 2–5 min. DNA was eluted in 30  $\mu\text{L}$  Milli-Q water and kept at room temperature for approximately 1 h. Extracted DNA was then stored at  $-20\text{ }^{\circ}\text{C}$  until further processing. All centrifugations were carried out at  $16,000\times g$  and  $4\text{ }^{\circ}\text{C}$  using a Model 157.MP RF centrifuge (Ole Dich).

The following real-time qPCR reaction mixture was used: 0.5  $\mu\text{L}$  of 10  $\mu\text{M}$  CDF670f (forward primer, 5'-GCAT CYTCTTGGWGAACG-3'), 0.5  $\mu\text{L}$  10  $\mu\text{M}$  CDF1058r (reverse primer, 5'-GTGCTGAAGGAGTCGT-3'), 1.5  $\mu\text{L}$   $\text{H}_2\text{O}$ , 5  $\mu\text{L}$  Brilliant II SYBR Green QRT-PCR Master Mix (Agilent Technologies) and 2.5  $\mu\text{L}$  sample. Samples (both environmental DNA and plasmids) were diluted 10- or 100-fold. The following qPCR program was used: initial soak at  $95\text{ }^{\circ}\text{C}$  for 10 min (following the manufacturer's protocol), 60 cycles of denaturation at  $95\text{ }^{\circ}\text{C}$  for 15 s, annealing at  $56\text{ }^{\circ}\text{C}$  for 15 s and extension at  $72\text{ }^{\circ}\text{C}$  for 30 s and, finally, melt curve analysis from 56 to  $95\text{ }^{\circ}\text{C}$  at  $0.5\text{ }^{\circ}\text{C}$  increments. Samples were run in triplicate and only those with a variation of  $<1\text{ Cq}$  were used. The plasmid from clone 010609\_08, a dinoflagellate phylotype from a previous study [37], was purified using the Cycle-Pure Kit (Omega Bio-Tek) and linearised [11] using the restriction enzyme NotI following the manufacturers' instructions and used as a standard in a 10-fold serial dilution. Real-time qPCR was performed on a CFX96 Touch Real-Time PCR Detection System (Bio-Rad). For the design of the dinoflagellate-specific primer pair CDF670f-CDF1058r, see [Supplementary material](#).

### Retention of Picoeukaryotes

To assess how well different filters retain picoeukaryotic organisms, their retention efficiency with respect to the smallest known eukaryote, *O. tauri* [6], was tested. *O. tauri* (RC1114) was obtained from the Roscoff Culture Collection and grown in L1 media [10] at  $20\text{ }^{\circ}\text{C}$  and 25  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ . A volume of 1 mL culture was filtered onto each of the following filters: GF/F (Whatman); 0.45- $\mu\text{m}$  standard MF mixed cellulose ester membrane (Millipore); 0.6- $\mu\text{m}$  polycarbonate filter (Poretics); and 0.8-, 0.4- and 0.2- $\mu\text{m}$  Nuclepore polycarbonate filter (Whatman). All filtrations were done at the same time from the same culture, which was thoroughly mixed prior to filtration. Except for the 0.2- $\mu\text{m}$  filtration, all filtrates were re-filtered on a 0.2- $\mu\text{m}$  filter, thus creating six size fractions: 0.2–0.8, 0.2–0.6, 0.2–0.45, 0.2–0.4, 0.2–GF/F and  $>0.2\text{ }\mu\text{m}$ . A drop of immersion oil on


both sides of the filter was used to attach the filter to the microscope slide and cover glass. An inverted microscope (Olympus IX81) equipped with a spinning disc system, epifluorescence and a digital camera (F-View III, Soft Imaging System) was used to take five images at random positions of each 0.2- $\mu\text{m}$  filter, and the number of cells was counted in each image. The 0.2- $\mu\text{m}$  filter was assumed to retain all *O. tauri* cells, and the relative retention of the other filters was calculated by comparing the >0.2- $\mu\text{m}$  size fraction with the pre-filtered size fractions.

## Results

### Literature Analysis

A total of 21 molecular picoeukaryotic libraries using universal eukaryotic primers were retrieved from the literature. Of these, only 15 met the criteria we had set for the analysis [1, 9, 17, 21, 25, 31, 32, 34, 36–38, 40] (Table S1), as three studies presented abundance in terms of operational taxonomical units rather than clones/tags [16, 24, 28] and another three used GF/F as end filters [3, 4, 29] (Table S2). A broad definition of picoeukaryotes is used in this analysis as the pre-filters used ranged from 1 to 5  $\mu\text{m}$  in pore size.

A Spearman's rank correlation found a significant negative relationship between the pore size of the end filter and the summed relative abundance of dinoflagellates and ciliates ( $r_s=0.63$ ,  $n=15$ ,  $p<0.02$ ) and an exponential regression explained  $r^2=0.51$  of the variation (Fig. 1). According to this regression, the abundance of these phylotypes decreases by 89 % when using a 0.8- $\mu\text{m}$  end filter as opposed to a 0.2- $\mu\text{m}$  end filter. Although the correlation is significant, considerable variation was noted for the results using 0.1–0.22- $\mu\text{m}$  end filters. No correlation was found between the abundance of rhizarians and the end filter used ( $r_s=0.13$ ,  $n=15$ ,  $p>0.50$ ), and the summed abundance of dinoflagellates and ciliates in



**Fig. 1** Summed proportion of dinoflagellates and ciliates observed, plotted against the pore size of the end filter used in published studies. There is a significant negative correlation and an exponential regression with an explained variation of  $r^2=0.51$

studies using GF/F (Table S2) did not differ significantly from studies using 0.2–0.22- $\mu\text{m}$  end filters (Table S1) in variance ( $F$  test,  $F=1.30$ ,  $df_1=2$ ,  $df_2=6$ ,  $p=0.68$ ) or mean ( $t$  test,  $t=0.26$ ,  $df=8$ ,  $p=0.80$ ).

### Real-Time qPCR


One sample from the 0.8–3- $\mu\text{m}$  size fraction was >20-fold higher than the average of the remaining four which only showed limited variation. This data point was identified as an outlier (Grubb's test,  $G=1.79$ ,  $p<0.01$ ) and excluded. The remaining data are shown in Fig. 2, where it can be seen that the amount of dinoflagellate 18S rDNA is significantly higher in the 0.2–3- $\mu\text{m}$  than in the 0.8–3- $\mu\text{m}$  size fraction ( $t$  test,  $t=2.64$ ,  $df=5$ ,  $p=0.046$ ) when data were log-transformed for variance homogeneity ( $F$  test,  $F=5.32$ ,  $df_1=2$ ,  $df_2=3$ ,  $p=0.21$ ). On average, the amount of dinoflagellate DNA decreased by 84 % when 0.8- $\mu\text{m}$  filters were used instead of the 0.2- $\mu\text{m}$  end filters, although the 0.2–3- $\mu\text{m}$  size fraction showed a considerable variation.

### Retention of Picoeukaryotes

On average, the 0.8- and 0.6- $\mu\text{m}$  polycarbonate filters retained 21 and 75 % of *O. tauri* cells, respectively. The remaining filters (0.4  $\mu\text{m}$  polycarbonate, 0.45  $\mu\text{m}$  mixed cellulose esters and GF/F) all retained 100 %.

## Discussion

The meta-analysis of molecular studies reporting the diversity of marine picoeukaryotes shows that the abundance of phylotypes belonging to ciliates and dinoflagellates decreases with increasing pore size of the end filter (Fig. 1). This suggests that either of these phylotypes



**Fig. 2** Relative dinoflagellate 18S rDNA copy number in two different size fractions as determined by real-time qPCR with standard deviations. The 0.2–3- $\mu\text{m}$  size fraction is set to 100 %. There is a significant difference between the two ( $t$  test,  $t=2.64$ ,  $df=5$ ,  $p=0.046$ ) when log transforming data for variance homogeneity ( $F$  test,  $F=5.32$ ,  $df_1=2$ ,  $df_2=3$ ,  $p=0.21$ )

originate from ciliates and dinoflagellates smaller than 1  $\mu\text{m}$  (and consequently not from organisms measuring 1–3  $\mu\text{m}$ ) or from extracellular DNA, which is more efficiently retained on filters with smaller rather than larger pore sizes. In light of the fact that 0.8  $\mu\text{m}$  is the smallest size ever recorded for eukaryotes [39] and the generally large (>10  $\mu\text{m}$ ) size of ciliates and dinoflagellates as well as the lack of microscopic observations of picoplanktonic ciliates and dinoflagellates, an extracellular origin of these phylotypes seems the most plausible explanation. Interestingly, the effect of end filter's pore size on the abundance of phylotypes of putative extracellular origin is nearly identical whether assessed by a meta-analysis of molecular studies of picoeukaryotes (Fig. 1) or experimentally using qPCR (Fig. 2). In both study approaches, the 0.2- $\mu\text{m}$  end filter shows a considerable variation and an average decrease of 84–89 % of phylotypes of suspected extracellular origin is observed when compared to the 0.8- $\mu\text{m}$  end filter. For the meta-analysis, the high variation for the 0.2- $\mu\text{m}$  end filters may be caused by a high general variation in ciliate and dinoflagellate abundance, while as a possible explanation for the variability with the qPCR approach, it can be noted that significant differences in the composition of the phytoplankton community have previously been detected at spatial scales much smaller than the resolution offered by the traditional Niskin bottles [30].

On this basis, it seems likely that reported observations of ciliates and dinoflagellates in marine picoeukaryotic molecular studies are filtration artefacts. This conclusion is supported by a study using fluorescence-assisted cell sorting (an approach that does not use sequential filtration, thereby avoiding potential filtration artefacts) to isolate the picoeukaryotic community from the English Channel, in that the study failed to find DNA sequences belonging to either group [20].

Dissolved extracellular DNA is ubiquitous in all marine environments [12] and can range from a few hundred to several thousand base pairs in length [8], fully encompassing the length of partial or complete 18S sequences. Furthermore, it can have a fast turnover time of mere hours [2, 19, 35]. Dissolved DNA is functionally defined as that which passes a 0.2- $\mu\text{m}$  filter [7, 12], but it can bind to particles in the marine sediment, such as sand, thereby inhibiting degradation of the DNA by DNase [14, 15] allowing for a much larger pool of extracellular DNA [5] and, possibly, also for retention on, for example, 0.2- $\mu\text{m}$  filters. Such particle-associated extracellular DNA has been shown to contain highly diverse 18S rDNA sequences [5] and could, possibly, be the source of both ciliate and dinoflagellate sequences in molecular picoeukaryotic studies.

In several molecular studies of picoeukaryotes, GF/F is used as the end filter (Table S2). Despite having a pore size of 0.7  $\mu\text{m}$  according to the manufacturer, GF/F in this study performed similarly to a 0.2- $\mu\text{m}$  filter, with retention of both

extracellular DNA and *O. tauri* cells. The fibrous nature of the filter is likely the cause of the apparent discrepancy between nominal pore size and retentive performance.

The largest decrease (84–89 %) in putative extracellular phylotypes compared to a 0.2- $\mu\text{m}$  end filter is seen when using a 0.8- $\mu\text{m}$  filter. Unfortunately, however, this pore size only retained 21 % of *O. tauri* cells in this study. For non-quantitative purposes, the 75 % retention of the 0.6- $\mu\text{m}$  filter might be sufficient but 0.4- $\mu\text{m}$  filters must be used if it is important that the entire picoeukaryotic community is retained. For investigations where only specifically larger picoeukaryotes are of interest (e.g. MAST clades), a pore size of 0.8  $\mu\text{m}$  might be prudent given the decreased retention of extracellular DNA compared to end filters with smaller pore sizes.

This study suggest that it is most likely that the rhizarian phylotypes recorded in the picoplanktonic fraction are not of extracellular origin as their abundance in molecular studies does not decrease with increasing pore size. This might be explained by the presence of rhizarian picoplanktonic swarmer cells which have been described [13].

The high abundance of phylotypes of a putative extracellular origin in molecular studies of picoeukaryotic diversity demonstrates the limitations of a purely molecular approach in describing marine plankton communities, as there is no clear link from phylotype to cellular identity. Visual confirmation (e.g. using flow cytometry or epifluorescence microscopy) of novel molecular diversity is necessary to demonstrate the existence of different phylotypes within the picoeukaryotic size fraction.

**Acknowledgments** The authors would like to thank Simon Bressendorff, Functional Genomics, Department of Biology, University of Copenhagen, for his very helpful assistance with real-time qPCR and the captain and crew of R/V Lance. The Fram Strait Cruise 2011 was arranged by the Norwegian Polar Institute. This work was supported by the Danish Council for Strategic Research (North Atlantic–Arctic coupling in a changing climate: impacts on ocean circulation, carbon cycling and sea-ice, 10-093003/DSF), Konto af 1988 for Botanisk Mikrobiologisk forskning, Center for Macroecology, Evolution and Climate supported by the Danish National Research Foundation and Department of Biology at the University of Copenhagen. ND thanks VILLUM FONDEN (VKR021561) for an equipment grant.

## References

1. Brown MV, Philip GK, Bunge JA, Smith MC, Bissett A, Lauro FM, Fuhrman JA, Donachie SP (2009) Microbial community structure in the North Pacific ocean. *ISME J* 3:1374–1386
2. Brum J (2005) Concentration, production and turnover of viruses and dissolved DNA pools at Stn ALOHA, North Pacific Subtropical Gyre. *Aquat Microb Ecol* 41:103–113
3. Cheung MK, Au CH, Chu KH, Kwan HS, Wong CK (2010) Composition and genetic diversity of picoeukaryotes in subtropical



- coastal waters as revealed by 454 pyrosequencing. *ISME J* 4:1053–1059
4. Cheung MK, Chu KH, Li CP, Kwan HS, Wong CK (2008) Genetic diversity of picoeukaryotes in a semi-enclosed harbour in the subtropical western Pacific Ocean. *Aquat Microb Ecol* 53:295–305
  5. Corinaldesi C, Barucca M, Luna GM, Dell'Anno A (2011) Preservation, origin and genetic imprint of extracellular DNA in permanently anoxic deep-sea sediments. *Mol Ecol* 20:642–654
  6. Courties C, Vaquer A, Troussellier M, Lautier J (1994) Smallest eukaryotic organism. *Nature* 370:255
  7. DeFlaun M, Paul J (1986) Simplified method for dissolved DNA determination in aquatic environments. *Appl Environ Microbiol* 52:654–659
  8. DeFlaun M, Paul J, Jeffrey W (1987) Distribution and molecular weight of dissolved DNA in subtropical estuarine and oceanic environments. *Mar Ecol Prog Ser* 38:65–73
  9. Díez B, Pedros-Alió C, Massana R (2001) Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing. *Appl Environ Microbiol* 67:2932–2941
  10. Guillard R, Hargraves P (1993) *Stichochrysis immobilis* is a diatom, not a chrysophyte. *Phycologia* 32:234–236
  11. Hou Y, Zhang H, Miranda L, Lin S (2010) Serious overestimation in quantitative PCR by circular (supercoiled) plasmid standard: microalgal *pcna* as the model gene. *PLoS One* 5:e9545
  12. Karl DM, Bailiff MD (2012) The measurement and distribution of dissolved nucleic acids in aquatic environments. *Limnol Oceanogr* 34:543–558
  13. Kimoto K, Yuasa T, Takahashi O (2011) Molecular identification of reproductive cells released from *Cypridopsis irregularis* Nigrini (Radiolaria). *Environ Microbiol Rep* 3:86–90
  14. Lorenz MG, Wackernagel W (1987) Adsorption of DNA to sand and variable degradation rates of adsorbed DNA. *Appl Environ Microbiol* 53:2948–2952
  15. Lorenz M, Aardema B (1981) Interaction of marine sediment with DNA and DNA availability to nucleases. *Mar Biol* 64:225–230
  16. Lovejoy C, Massana R, Pedros-Alió C, Pedro C (2006) Diversity and distribution of marine microbial eukaryotes in the Arctic Ocean and adjacent seas. *Appl Environ Microbiol* 72:3085–3095
  17. Lovejoy C, Potvin M (2011) Microbial eukaryotic distribution in a dynamic Beaufort Sea and the Arctic Ocean. *J Plankton Res* 33:431–444
  18. López-García P, Rodríguez-Valera F, Pedros-Alió C, Moreira D (2001) Unexpected diversity of small eukaryotes in deep-sea. Antarctic plankton *Nature* 409:603–607
  19. Løvdal T, Tanaka T, Thingstad TF (2007) Algal–bacterial competition for phosphorus from dissolved DNA, ATP, orthophosphate in a mesocosm experiment. *Limnol Oceanogr* 52:1407–1419
  20. Marie D, Shi XL, Rigaut-Jalabert F, Vault D (2010) Use of flow cytometric sorting to better assess the diversity of small photosynthetic eukaryotes in the English Channel. *FEMS Microbiol Ecol* 72:165–178
  21. Marie D, Zhu F, Balagué V, Ras J, Vault D (2006) Eukaryotic picoplankton communities of the Mediterranean Sea in summer assessed by molecular approaches (DGGE, TTGE, QPCR). *FEMS Microbiol Ecol* 55:403–415
  22. Massana R, Castresana J, Balagué V, Guillou L, Romari K, Groisillier A, Valentin K, Pedros-Alió C (2004) Phylogenetic and ecological analysis of novel marine stramenopiles. *Appl Environ Microbiol* 70:3528–3534
  23. Massana R (2011) Eukaryotic picoplankton in surface oceans. *Annu Rev Microbiol* 65:91–110
  24. Massana R, Balagué V, Guillou L, Pedros-Alió C (2004) Picoeukaryotic diversity in an oligotrophic coastal site studied by molecular and culturing approaches. *FEMS Microbiol Ecol* 50:231–243
  25. Massana R, Guillou L, Terrado R, Forn I, Pedros-Alió C (2006) Growth of uncultured heterotrophic flagellates in unamended sea-water incubations. *Aquat Microb Ecol* 45:171–180
  26. Massana R, Pedros-Alió C (2008) Unveiling new microbial eukaryotes in the surface ocean. *Curr Opin Microbiol* 11:213–218
  27. Massana R, Terrado R, Forn I, Lovejoy C, Pedros-Alió C (2006) Distribution and abundance of uncultured heterotrophic flagellates in the world oceans. *Environ Microbiol* 8:1515–1522
  28. Medlin LK, Metfies K, Mehl H, Wiltshire K, Valentin K (2006) Picoeukaryotic plankton diversity at the Helgoland time series site as assessed by three molecular methods. *Microb Ecol* 52:53–71
  29. Moon-van der Staay SY, De Wachter R, Vault D (2001) Oceanic 18S rDNA sequences from picoplankton reveal unsuspected eukaryotic diversity. *Nature* 409:607–610
  30. Mouritsen LT, Richardson K (2003) Vertical microscale patchiness in nano- and microplankton distributions in a stratified estuary. *J Plankton Res* 25:783–797
  31. Not F, Gausling R, Azam F, Heidelberg JF, Worden AZ (2007) Vertical distribution of picoeukaryotic diversity in the Sargasso Sea. *Environ Microbiol* 9:1233–1252
  32. Not F, Latasa M, Scharek R, Viprey M, Karleskind P, Balagué V, Ontoria-Oviedo I, Cumino A, Goetze E, Vault D, Massana R (2008) Protistan assemblages across the Indian Ocean, with a specific emphasis on the picoeukaryotes. *Deep-Sea Res I Oceanogr Res Pap* 55:1456–1473
  33. Not F, Valentin K, Romari K, Lovejoy C, Massana R, Töbe K, Vault D, Medlin LK (2007) Picobiphilophytes: a marine picoplanktonic algal group with unknown affinities to other eukaryotes. *Science* 315:253–255
  34. Not F, del Campo J, Balagué V, de Vargas C, Massana R (2009) New insights into the diversity of marine picoeukaryotes. *PLoS One* 4:e7143
  35. Paul JH, Jeffrey WH, DeFlaun MF (1987) Dynamics of extracellular DNA in the marine environment. *Appl Environ Microbiol* 53:170–179
  36. Romari K, Vault D (2004) Composition and temporal variability of picoeukaryote communities at a coastal site of the English Channel from 18S rDNA sequences. *Limnol Oceanogr* 49:784–798
  37. Sørensen N, Daugbjerg N, Gabrielsen TM (2011) Molecular diversity and temporal variation of picoeukaryotes in two Arctic fjords, Svalbard. *Polar Biol* 35:519–533
  38. Terrado R, Vincent WF, Lovejoy C (2009) Mesopelagic protists: diversity and succession in a coastal Arctic ecosystem. *Aquat Microb Ecol* 56:25–39
  39. Vault D, Eikrem W, Viprey M, Moreau H (2008) The diversity of small eukaryotic phytoplankton ( $\leq 3 \mu\text{m}$ ) in marine ecosystems. *FEMS Microbiol Rev* 32:795–820
  40. Worden A (2006) Picoeukaryote diversity in coastal waters of the Pacific Ocean. *Aquat Microb Ecol* 43:165–175